

THE PROBABILITY OF CHOOSING PRIMITIVE SETS

SERGI ELIZALDE AND KEVIN WOODS

ABSTRACT. We generalize a theorem of Nymann that the density of points in \mathbb{Z}^d that are visible from the origin is $1/\zeta(d)$, where $\zeta(a)$ is the Riemann zeta function $\sum_{i=1}^{\infty} 1/i^a$. A subset $S \subset \mathbb{Z}^d$ is called primitive if it is a \mathbb{Z} -basis for the lattice $\mathbb{Z}^d \cap \text{span}_{\mathbb{R}}(S)$, or, equivalently, if S can be completed to a \mathbb{Z} -basis of \mathbb{Z}^d . We prove that if m points in \mathbb{Z}^d are chosen uniformly and independently at random from a large box, then as the size of the box goes to infinity, the probability that the points form a primitive set approaches $1/(\zeta(d)\zeta(d-1)\cdots\zeta(d-m+1))$.

1. INTRODUCTION

A classic result in number theory is that, if a point in \mathbb{Z}^2 is chosen “at random,” the probability that the point is visible from the origin (that is, not hidden by another point in \mathbb{Z}^2) is $\frac{1}{\zeta(2)}$, where $\zeta(a)$ is the Riemann zeta function $\sum_{i=1}^{\infty} \frac{1}{i^a}$ (see [1] for a proof using Euler’s totient function). More precisely, for a given n , if we choose an integer point (a, b) uniformly at random from the box $[-n, n] \times [-n, n]$ and compute the probability that (a, b) is visible from the origin, then as n approaches infinity, this probability approaches $\frac{1}{\zeta(2)}$.

J.E. Nymann generalized this result to higher dimensions [7]: if a point in \mathbb{Z}^d is chosen at random, then the probability that the point is visible from the origin is $\frac{1}{\zeta(d)}$. This theorem is true for $d \geq 2$ and is, in effect, true for $d = 1$: the only points in \mathbb{Z}^1 that are visible from the origin are ± 1 , so the probability is zero, and $\zeta(1)$ diverges so that $\frac{1}{\zeta(1)} = 0$.

An obvious way to restate the condition that a point $s = (a_1, a_2, \dots, a_d) \in \mathbb{Z}^d$ is visible from the origin is that $\gcd(a_1, \dots, a_d) = 1$. We will restate the condition in a lattice theoretic context, so that it may be generalized to picking more than one point in \mathbb{Z}^d . A point s is visible from the origin if and only if $\{s\}$ is a \mathbb{Z} -basis for the lattice $\text{span}_{\mathbb{R}}(s) \cap \mathbb{Z}^d$. In general, given a set $S = \{s_1, s_2, \dots, s_m\} \subset \mathbb{Z}^d$, where $1 \leq m \leq d$, we say that S is *primitive* if S is a \mathbb{Z} -basis for the lattice $\text{span}_{\mathbb{R}}(S) \cap \mathbb{Z}^d$. An equivalent definition [6] is that S is primitive if and only if S can be completed to a \mathbb{Z} -basis of all of \mathbb{Z}^d .

In this paper we prove that if S is chosen “at random,” then the probability that S is primitive is

$$\frac{1}{\zeta(d)\zeta(d-1)\cdots\zeta(d-m+1)}.$$

To be precise, we prove the following theorem.

Theorem 1. *Let d and m be given, with $m \leq d$. For $n \in \mathbb{Z}_+$, $1 \leq k \leq m$, and $1 \leq i \leq d$, let $b_{n,k,i} \in \mathbb{Z}$. For a given n , choose integers s_{ki} uniformly (and independently) at random from the set $b_{n,k,i} \leq s_{ki} < b_{n,k,i} + n$. Let $s_k = (s_{k1}, \dots, s_{kd})$ and let $S = \{s_1, s_2, \dots, s_m\}$.*

If $m < d$ and $|b_{n,k,i}|$ is bounded by a polynomial in n , then, as n approaches infinity, the probability that S is a primitive set approaches

$$\frac{1}{\zeta(d)\zeta(d-1)\cdots\zeta(d-m+1)},$$

where $\zeta(a)$ is the Riemann zeta function $\sum_{i=1}^{\infty} \frac{1}{i^a}$.

If $m = d$, then, as n approaches infinity, the probability that S is a primitive set approaches zero.

When $m = 1$, this theorem gives the classic result ($d = 2$) and Nymann’s result. The statements for $m < d$ and $m = d$ are consistent, because for $m = d$ the value of

$$\frac{1}{\zeta(d)\zeta(d-1)\cdots\zeta(1)}$$

is zero in the sense that $\zeta(1)$ diverges.

The statement of the theorem uses more general boxes than $[-n, n]^d$ to pick the s_k from. We do this because the more general result is needed in [4]. That paper was the original inspiration for this theorem: we discovered the theorem in an attempt to prove a fact in computational biology and Bayesian network theory. Since the concept of primitive sets is important in the geometry of numbers, we are proving this theorem in this separate paper.

Note that some bound on the $b_{n,k,i}$ in terms of n is needed; otherwise one could construct arbitrarily large boxes from which *no* primitive sets could be selected (even for $d = 2$, $m = 1$), as the following proposition shows.

Proposition 2. *Given $n \in \mathbb{Z}_+$, there exists integers $b_{n,1,1}$ and $b_{n,1,2}$ such that no integer vectors (s_{11}, s_{12}) chosen from the box*

$$b_{n,1,1} \leq s_{11} < b_{n,1,1} + n \text{ and } b_{n,1,2} \leq s_{12} < b_{n,1,2} + n$$

are visible from the origin.

Proof. Given n , choose n^2 distinct primes, p_{ij} , for $0 \leq i, j < n$. For $0 \leq i < n$, let $P_i = \prod_{j=0}^{n-1} p_{ij}$. For $0 \leq j < n$, let $Q_j = \prod_{i=0}^{n-1} p_{ij}$. Since the P_i are relatively prime, we may use the Chinese Remainder Theorem to choose a $b_{n,1,1}$ such that

$$b_{n,1,1} \equiv -i \pmod{P_i}, \text{ for } 0 \leq i < n,$$

in other words, so that P_i divides $b_{n,1,1} + i$. Similarly, choose a $b_{n,1,2}$ so that Q_j divides $b_{n,1,2} + j$, for $0 \leq j < n$. Then for any choice of $(s_{11}, s_{12}) = (b_{n,1,1} + i, b_{n,1,2} + j)$ from the box, p_{ij} divides s_{11} and s_{12} , and the point is not visible from the origin. \square

In Section 2, we present an outline of the proof of Theorem 1. The outline is a full proof in every respect, except that we ignore the error estimations in our probabilities. In that sense, it is the “moral” proof of the result. In Section 3, we fill the holes by proving that the error estimates approach zero as n approaches infinity. The methods in Section 3 are themselves of interest, using concepts from triangulations of point sets, the metric geometry of polytopes (cross-sections of d-cubes), analytic number theory (consequences of the Prime Number Theorem), and the geometry of numbers.

2. OUTLINE OF THE PROOF

We first prove the more difficult case $m < d$. At the end of this section, we will prove the $m = d$ case. We proceed by induction on m .

If $m = 0$, the theorem is trivially true. Assume that the theorem is true for $m - 1$, and we will prove it for m . The probability that $S = \{s_1, s_2, \dots, s_m\}$ is primitive is the product

$$\text{Prob}_{\mathcal{P}_n}(\{s_1, \dots, s_{m-1}\} \text{ is primitive}) \\ \cdot \text{Prob}_{\mathcal{P}_n}(S \text{ is primitive, given that } \{s_1, \dots, s_{m-1}\} \text{ is primitive}),$$

where \mathcal{P}_n is the probability distribution, for a given n , from which we are choosing S . The first term in the product approaches

$$\frac{1}{\zeta(d)\zeta(d-1)\cdots\zeta(d-m+2)},$$

as $n \rightarrow \infty$, by the inductive hypothesis, so we must show that the second term approaches $\frac{1}{\zeta(d-m+1)}$.

Indeed, suppose $\{s_1, \dots, s_{m-1}\}$ is given and is primitive, and we choose $s_m = (s_{m1}, \dots, s_{md})$ (independently from the other s_i) according to the probability distribution \mathcal{P}_n . Let A be the $(m-1) \times d$ integer matrix whose rows are s_1, \dots, s_{m-1} . We will need the following lemma, to find a simpler matrix whose rows also form a primitive set.

Lemma 3. *Let A be a matrix in $\mathbb{Z}^{p \times q}$, and let U be a unimodular matrix (i.e., $\det(U) = \pm 1$) in $\mathbb{Z}^{q \times q}$. The rows of A form a primitive set if and only if the rows of AU also form a primitive set.*

Proof. Suppose the rows of A form a primitive set. Let $a \in \mathbb{Z}^q$ be in the \mathbb{R} -span of the rows of AU , that is, $a = xAU$, where x is a matrix in $\mathbb{R}^{1 \times p}$. In order to show that the rows of AU form a primitive set, we must show that x is actually integral. Indeed, $aU^{-1} = xA \in \mathbb{Z}^q$ is in the \mathbb{R} -span of the rows of A , and since the rows of A form a primitive set, x must be integral. This also proves the converse, as U^{-1} is unimodular and $A = (AU)U^{-1}$. \square

The matrix U we will choose is a matrix that puts AU into *Hermite normal form*.

Definition 4. A matrix $B \in \mathbb{Z}^{p \times q}$ is in Hermite normal form if

- (1) $B_{ij} = 0$ for all $j > i$,
- (2) $B_{ii} > 0$ for all i , and
- (3) $0 \leq B_{ij} < B_{ii}$ for all $j < i$.

Given any integer matrix B of full row rank, there exists a unimodular matrix U such that BU is in Hermite normal form (see, e.g., [5]; U will not, in general, be unique). This fact, together with the following lemma, gives a convenient characterization of when S is a primitive set.

Lemma 5. Let $\{s_1, \dots, s_{m-1}\} \subset \mathbb{Z}^d$ be a primitive set, and let $s_m \in \mathbb{Z}^d$ be given. Let A be the (full row rank) matrix with rows s_1, \dots, s_{m-1} , and let U be a matrix such that AU is in Hermite normal form. Let $U^{(i)}$ be the i -th column of U . Then $\{s_1, \dots, s_m\}$ is a primitive set if and only if the $s_m U^{(i)}$, for $m \leq i \leq d$, are relatively prime.

Proof. By Lemma 3, the rows of AU form a primitive set. It follows that $(AU)_{ii} = 1$, for $1 \leq i \leq m-1$ (otherwise e_i , the i -th standard basis vector, would be in the \mathbb{R} -span of the rows of AU , but not in the \mathbb{Z} -span). Then, from the definition of Hermite normal form, $(AU)_{ij} = 0$ for $i \neq j$. Let A' be the matrix with rows s_1, \dots, s_m (that is, A' is A with the additional row s_m appended). By Lemma 3, $\{s_1, \dots, s_m\}$ is a primitive set if and only if the rows of $A'U$ form a primitive set. We see that this is true if and only if the $(A'U)_{mi}$, for $m \leq i \leq d$, are relatively prime (indeed, the index of the lattice $\text{span}_{\mathbb{Z}}\{s_1, \dots, s_m\}$ within $\mathbb{Z}^d \cap \text{span}_{\mathbb{R}}\{s_1, \dots, s_m\}$ is $\gcd\{(A'U)_{mi} : m \leq i \leq d\}$). Since $(A'U)_{mi} = s_m U^{(i)}$, the lemma follows. \square

Let $\mu : \mathbb{Z}_+ \rightarrow \{-1, 0, 1\}$ be the Möbius function defined to be

$$\mu(D) = \begin{cases} (-1)^i & \text{if } D \text{ is the product of } i \text{ distinct primes,} \\ 0 & \text{if } D \text{ is divisible by the square of a prime.} \end{cases}$$

Given $D \in \mathbb{Z}_+$, let p_{nD} be the probability that D divides $s_m U^{(i)}$ for all $m \leq i \leq d$. Note that p_{nD} is independent of our choice of U , because, as we noted in the proof of Lemma 5, $\gcd\{s_m U^{(i)} : m \leq i \leq d\}$ is the index of the lattice $\text{span}_{\mathbb{Z}}\{s_1, \dots, s_m\}$ within $\mathbb{Z}^d \cap \text{span}_{\mathbb{R}}\{s_1, \dots, s_m\}$, which is independent of U . Then, using inclusion-exclusion, the probability that the $s_m U^{(i)}$, for $m \leq i \leq d$, are relatively prime is

$$\sum_{D=1}^{\infty} \mu(D) p_{nD}.$$

We expect each p_{nD} to be approximately $D^{-(d-m+1)}$. In Section 3, we will show that

$$(1) \quad \lim_{n \rightarrow \infty} \sum_{D=1}^{\infty} \mu(D) p_{nD} = \sum_{D=1}^{\infty} \mu(D) D^{-(d-m+1)}.$$

Given that we have verified (1), the following well-known lemma (see [1, Section 11.4] for a proof), applied to $a = d - m + 1$, finishes the proof of the theorem for the $m < d$ case.

Lemma 6. *For any integer $a \geq 2$,*

$$\sum_{D=1}^{\infty} \mu(D) D^{-a} = \frac{1}{\zeta(a)}.$$

To conclude this section, we prove the $m = d$ case. Suppose we have chosen $S' = \{s_1, s_2, \dots, s_{d-1}\}$. If S' is not primitive, then there is no choice of s_d that will make the full set $S = \{s_1, s_2, \dots, s_d\}$ primitive. If S' is primitive, then consider the hyperplane $W = \text{span}_{\mathbb{R}}(S')$. Choose a vector $a \in \mathbb{Z}_d$, whose coordinates are relatively prime, such that

$$W = \{x \in \mathbb{R}^d : \langle a, x \rangle = 0\}$$

(where $\langle \cdot, \cdot \rangle$ is the standard dot product).

Then S will be primitive (a \mathbb{Z} -basis for \mathbb{Z}^d) if and only if $\langle a, s_d \rangle = \pm 1$. As n approaches infinity, we would expect, of the n^d possible choices for s_d , the number that lie on these two hyperplanes to be $O(n^{d-1})$. And indeed it is: the precise error estimation follows by Claim 2a' in the proof of Lemma 9.

Since the number of possible choices for s_d is n^d and the number that make S primitive is $O(n^{d-1})$, the probability that S is primitive approaches zero.

3. ERROR ESTIMATES

The remaining piece of the proof, in the $m < d$ case, is to demonstrate Equation (1), that is, that

$$\left| \sum_{D=1}^{\infty} \mu(D) p_{nD} - \sum_{D=1}^{\infty} \mu(D) D^{-(d-m+1)} \right| \rightarrow 0$$

as $n \rightarrow \infty$.

We will need a bound on the entries of U , which the following lemma will help us get.

Lemma 7. *Given a matrix $A \in \mathbb{Z}^{p \times q}$ of full row rank and a bound M_0 such that $|A_{ij}| < M_0$ for all i, j , there exists a unimodular matrix U such that*

- (1) AU is in Hermite normal form and
- (2) $|U_{ij}| \leq p!qM_0^p$ for all i, j .

Proof. Let B be the $q \times q$ matrix obtained by appending to A the rows e_1, e_2, \dots, e_{q-p} (where e_i is the i -th standard basis vector). Without loss of generality, we can assume that B is a nonsingular matrix (otherwise, we could have appended different e_i). Let U be a unimodular matrix such that BU is in Hermite normal form. Note that AU is also in Hermite normal form.

We will use the fact that

$$(2) \quad U = B^{-1}(BU) = \frac{1}{\det(B)} \operatorname{adj}(B)(BU),$$

where $\operatorname{adj}(B)$ is the adjugate (classical adjoint) of B , in order to bound the entries of U . Since BU is lower diagonal,

$$|\det(B)| = \det(BU) = \prod_{i=1}^q (BU)_{ii}.$$

Therefore $(BU)_{ii} \leq |\det(B)|$ for all i , and, by the definition of Hermite normal form, we conclude that $(BU)_{ij} \leq |\det(B)|$ for all i, j .

Since the first p rows of B have entries bounded by M_0 and the remaining rows are standard basis vectors, the entries of $\operatorname{adj}(B)$ are bounded by $p!M_0^p$. Combining these two bounds, we see that the entries of $\operatorname{adj}(B)(BU)$ are bounded by $q \cdot p!M_0^p \cdot |\det(B)|$. Using (2) we conclude that

$$|U_{ij}| \leq \frac{1}{|\det(B)|} q \cdot p!M_0^p \cdot |\det(B)| = p!qM_0^p$$

for all i, j , as desired. □

Since the absolute value of the entries of A are bounded by $|b_{n,k,i}| + n$, which we assume to be bounded by a polynomial in n , Lemma 7 shows that the unimodular matrix U can be chosen such that the absolute value of each entry of U is bounded by a polynomial in n . This in turn implies that $|s_m U^{(i)}|$ is also bounded by a polynomial in n (where $U^{(i)}$ is the i -th column of U). Let $M = M(n)$ be our bound on $|s_m U^{(i)}|$; say M is $O(n^k)$ for some k . Clearly, for $D > M$, $p_{nD} = 0$.

We have that

$$\begin{aligned}
(3) \quad & \left| \sum_{D=1}^{\infty} \mu(D) p_{nD} - \sum_{D=1}^{\infty} \mu(D) D^{-(d-m+1)} \right| \\
& \leq \left| \sum_{D=1}^n \mu(D) (p_{nD} - D^{-(d-m+1)}) \right| + \left| \sum_{D=n+1}^M \mu(D) p_{nD} \right| \\
& \quad + \left| \sum_{D=M+1}^{\infty} \mu(D) p_{nD} \right| + \left| \sum_{D=n+1}^{\infty} \mu(D) D^{-(d-m+1)} \right| \\
& \leq \sum_{D=1}^n |p_{nD} - D^{-(d-m+1)}| + \sum_{D=n+1}^M p_{nD} + 0 + \sum_{D=n+1}^{\infty} D^{-(d-m+1)}.
\end{aligned}$$

Of the three nonzero terms in the last expression, $\sum_{D=n+1}^{\infty} D^{-(d-m+1)}$ certainly converges to zero as n approaches infinity, so it suffices to show that the first two terms, $\sum_{D=1}^n |p_{nD} - D^{-(d-m+1)}|$ and $\sum_{D=n+1}^M p_{nD}$, do as well. We break our error computation into these two cases.

Before we handle the two error sums in Lemmas 8 and 9, we set some common terminology. Let \mathcal{B}_n be the d -dimensional box of integers $\{s_m \in \mathbb{Z}^d : b_{n,m,i} \leq s_{mi} < b_{n,m,i} + n, \text{ for all } i\}$, which is the box from which s_m is chosen with uniform probability. Given $D \in \mathbb{Z}_+$, let $\Lambda_D \subset \mathbb{Z}^d$ be the lattice of integer vectors $x \in \mathbb{Z}^d$ such that D divides $x \cdot U^{(i)}$, for $m \leq i \leq d$. Λ_D is a sublattice of \mathbb{Z}^d of index D^{d-m+1} . Let $S_{nD} = \mathcal{B}_n \cap \Lambda_D$. Then

$$(4) \quad p_{nD} = \frac{|S_{nD}|}{n^d}.$$

Lemma 8. *As defined above,*

$$\sum_{D=1}^n |p_{nD} - D^{-(d-m+1)}|$$

converges to zero as $n \rightarrow \infty$.

Proof. Suppose $1 \leq D \leq n$. Let $L_D \subset \mathbb{Z}^d$ be the lattice of integer vectors $(x_1, \dots, x_d) \in \mathbb{Z}^d$ such that D divides each x_i . L_D is a sublattice of \mathbb{Z}^d of index D^d . In fact, we see that L_D is a sublattice of Λ_D , and therefore its index in Λ_D is $D^d / D^{d-m+1} = D^{m-1}$.

This means that if we look at any $D \times \dots \times D$ cube, $C = \{(x_1, \dots, x_d) \in \mathbb{Z}^d : r_i \leq x_i < r_i + D\}$ for some $r_i \in \mathbb{Z}$ (that is, a translate of a fundamental parallelepiped of L_D), then C contains exactly D^{m-1} elements of Λ_D . Since \mathcal{B}_n can be covered by $(\frac{n}{D} + 1)^d$ such boxes, we have that $|S_{nD}| \leq D^{m-1} (\frac{n}{D} + 1)^d$, and so

$$p_{nD} \leq \frac{D^{m-1} (\frac{n}{D} + 1)^d}{n^d} = D^{m-1-d} \left(1 + \frac{D}{n}\right)^d.$$

Similarly, $(\frac{n}{D} - 1)^d$ disjoint $D \times \cdots \times D$ cubes can be placed inside \mathcal{B}_n , and so

$$p_{nD} \geq D^{m-1-d} \left(1 - \frac{D}{n}\right)^d.$$

Combining these two inequalities, for some c with $|c| \leq 1$ we have that

$$p_{nD} = D^{m-1-d} \left(1 + c \frac{D}{n}\right)^d = D^{m-1-d} \left(1 + O\left(\frac{dD}{n}\right)\right).$$

It follows that

$$\left|p_{nD} - \frac{1}{D^{d-m+1}}\right| \leq D^{m-d} O\left(\frac{d}{n}\right)$$

and so

$$\sum_{D=1}^n \left|p_{nD} - D^{-(d-m+1)}\right| \leq O\left(\frac{d}{n}\right) \sum_{D=1}^n D^{m-d},$$

which converges to zero as $n \rightarrow \infty$, proving the lemma. \square

Lemma 9. *As defined above,*

$$\sum_{D=n+1}^M p_{nD},$$

converges to zero as $n \rightarrow \infty$.

Proof. Let

$$T_n = \bigcup_{D=n+1}^M S_{nD}.$$

Let N_n be the maximum, over all $s_m \in \mathcal{B}_n$, of

$$\#\{D : n < D \leq M \text{ and } s_m \in S_{nD}\}.$$

Then

$$\begin{aligned} \sum_{D=n+1}^M p_{nD} &= n^{-d} \sum_{D=n+1}^M |S_{nD}| \\ &\leq n^{-d} |T_n| \cdot N_n \end{aligned}$$

We need to approximate N_n and $|T_n|$. We will repeatedly use the following fact (see [1], p:294), which can be derived from the Prime Number Theorem: for any $\epsilon > 0$ and for any $r \leq M$, the number of factors of r is $O(n^\epsilon)$ (more precisely, for any $\delta > 0$ and sufficiently large r , the number of factors of r is less than $r^{(1+\delta) \log 2 / \log \log r}$; now we use that $r \leq M$ is $O(n^k)$ for some k).

Claim 1: N_n is $O(n^\epsilon)$.

This follows immediately, as any element of the set

$$\{D : n < D \leq M \text{ and } s_m \in S_{nD}\}$$

must be a factor of, say, $s_m U^{(m)}$, and this number has $O(n^\epsilon)$ factors.

Claim 2: $|T_n|$ is $O(n^{d-\frac{1}{2}+\epsilon})$.

Let $a = \gcd(U_1^{(i)} : m \leq i \leq d)$, where $U^{(m)}, U^{(m+1)}, \dots, U^{(d)}$ are the last $d-m+1$ columns of U . Let R be the set of integers greater than n that are factors of at least one of $a, 2a, 3a, \dots, \lfloor \sqrt{n} \rfloor a$. Each of the $\lfloor \sqrt{n} \rfloor$ numbers $i \cdot a$ such that $1 \leq i \leq \lfloor \sqrt{n} \rfloor$ has $O(n^\epsilon)$ factors, so $|R|$ is $O(n^{\frac{1}{2}+\epsilon})$.

We divide T_n into two parts. Let

$$T_{n1} = \bigcup_{D \in R} S_{nD}$$

and let $T_{n2} = T_n \setminus T_{n1}$. We will show that both $|T_{n1}|$ and $|T_{n2}|$ are $O(n^{d-\frac{1}{2}+\epsilon})$, and so it will follow that $|T_n| = |T_{n1}| + |T_{n2}|$ is also $O(n^{d-\frac{1}{2}+\epsilon})$.

Claim 2a: $|T_{n1}|$ is $O(n^{d-\frac{1}{2}+\epsilon})$.

Given a $D \in R$, we want to estimate how large S_{nD} is. Suppose first that $\text{conv}(S_{nD})$ is a full dimensional polytope in \mathbb{Z}^d , that is, its affine hull is all of \mathbb{R}^d . Triangulate $\text{conv}(S_{nD})$ into at least $|S_{nD}| - d$ simplices whose vertices are in S_{nD} (this can always be done, see for example [3]). Each simplex in the triangulation has volume at least $\frac{1}{d!} D^{d-m+1}$, because the lattice Λ_n (which includes every point in S_{nD}) has index D^{d-m+1} in \mathbb{Z}^d . But $\text{conv}(S_{nD})$ has volume at most n^d , because it lies in \mathcal{B}_n . Putting this together,

$$\frac{1}{d!} D^{d-m+1} (|S_{nD}| - d) \leq n^d,$$

and so

$$|S_{nD}| \leq d + d! \frac{n^d}{D^{d-m+1}} \leq d + d! n^{m-1},$$

which is $O(n^{m-1})$.

On the other hand, if $\text{conv}(S_{nD})$ is not full dimensional, then Claim 2a', following, demonstrates that $|S_{nD}|$ is $O(n^{d-1})$. Therefore, in either case, $|S_{nD}|$ is $O(n^{d-1})$, and since $|R|$ is $O(n^{\frac{1}{2}+\epsilon})$, $|T_{n1}|$ is $O(n^{d-1} \cdot n^{\frac{1}{2}+\epsilon}) = O(n^{d-\frac{1}{2}+\epsilon})$, and Claim 2a follows.

Claim 2a': Let $X \subset \mathbb{Z}^d \cap \mathcal{B}_n$ (where \mathcal{B}_n is the box we are choosing s_m from). If the affine hull of X is k -dimensional, then $|X|$ is $O(n^k)$.

Let H be the k -dimensional affine space such that $X \subset H$. The k -dimensional Euclidean volume of $H \cap \mathcal{B}_n$ is at most $\sqrt{2}^{d-k} n^k$, as proved in [2]. Again we can triangulate $\text{conv}(X)$ into at least $|X| - k$ simplices that are k -dimensional. The best we can know this time is that each simplex has volume at least $\frac{1}{k!}$. Putting this together,

$$\frac{1}{k!}(|X| - k) \leq \sqrt{2}^{d-k} n^k,$$

and so $|X|$ is $O(n^k)$, proving Claim 2a'.

Claim 2b: $|T_{n2}|$ is $O(n^{d-\frac{1}{2}+\epsilon})$.

Recall that $a = \gcd(U_1^{(i)} : m \leq i \leq d)$. Without loss of generality, we may assume that $U_1^{(m)} = a$ and $U_1^{(i)} = 0$, for $m+1 \leq i \leq d$ (if not, we may perform elementary column operations on the last $d-m+1$ columns of U in order to put them in that form; the matrix AU will remain in Hermite normal form, because the last $d-m+1$ columns of AU are all zeros). Note that $a < M$.

Now suppose $s_{m2}, s_{m3}, \dots, s_{md}$ are given, such that $b_{n,m,i} \leq s_{mi} < b_{n,m,i} + n$. Given j such that $b_{n,m,1} \leq j < b_{n,m,1} + n$, define

$$t^{(j)} = (j, s_{m2}, s_{m3}, \dots, s_{md}).$$

We will show that $O(n^{\frac{1}{2}+\epsilon})$ of the $t^{(j)}$ are in T_{n2} (for given s_{m2}, \dots, s_{md}).

Since $U_1^{(m+1)} = 0$, $s' := t^{(j)}U^{(m+1)}$ is independent of j . If $t^{(j)} \in S_{nD}$ for a particular D , then D must be a factor of s' , which has $O(n^\epsilon)$ factors. Therefore there are only $O(n^\epsilon)$ possible D for which any of the $t^{(j)}$ could be a member of S_{nD} .

Now let us consider, for a given $D \notin R$, how many of the $t^{(j)}$ could be in S_{nD} . If $t^{(j)}$ and $t^{(k)}$ are in S_{nD} , then D divides $t^{(j)}U^{(m)}$ and $t^{(k)}U^{(m)}$. Therefore D divides the difference $t^{(j)}U^{(m)} - t^{(k)}U^{(m)}$, which is $(j-k) \cdot a$, since $U_1^{(m)} = a$. Since $D \notin R$, D does not divide $a, 2a, \dots, \lfloor \sqrt{n} \rfloor a$, and so $|j-k| > \sqrt{n}$. Therefore the number of j such that $t^{(j)} \in S_{nD}$ is at most $n/\sqrt{n} = \sqrt{n}$.

Since there are $O(n^\epsilon)$ possibilities for D , and since, for a given $D \notin R$, the number of $t^{(j)}$ in S_{nD} is $O(n^{\frac{1}{2}})$, we conclude that $O(n^{\frac{1}{2}+\epsilon})$ of the $t^{(j)}$ are in T_{n2} .

Since there are n^{d-1} choices for s_{m2}, \dots, s_{md} , we have that $|T_{n2}|$ is

$$O(n^{d-1} n^{\frac{1}{2}+\epsilon}) = O(n^{d-\frac{1}{2}+\epsilon}),$$

proving Claim 2b.

Combining our estimates of N_n and $|T_n|$ from Claims 1 and 2, we have that

$$\begin{aligned} \sum_{D=n+1}^M p_{nD} &\leq n^{-d} |T_n| \cdot N_n \\ &= n^{-d} O(n^{d-\frac{1}{2}+\epsilon}) O(n^\epsilon) \\ &= O(n^{-\frac{1}{2}+2\epsilon}), \end{aligned}$$

and therefore $\sum_{D=n+1}^M p_{nD}$ converges to zero as n approaches infinity. \square

Combining Lemmas 8 and 9 with Equation (3), we have shown that

$$\left| \sum_{D=1}^{\infty} \mu(D) p_{nD} - \sum_{D=1}^{\infty} D^{-(d-m+1)} \right| \rightarrow 0$$

as $n \rightarrow \infty$. This completes our error analysis and, together with Section 2, provides a complete proof of Theorem 1.

Acknowledgements. The authors are grateful to Carl Pomerance for several helpful discussions and to Ravi Kannan for the proof of Lemma 7.

REFERENCES

- [1] T.M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, New York (1976).
- [2] K. Ball, Volumes of sections of cubes and related problems. *Geometric aspects of functional analysis (1987-88)*, Lecture Notes in Mathematics, Springer, Berlin (1989), 251–260.
- [3] J.A. De Loera, J. Rambau, and F. Santos Leal, *Triangulations: Applications, Structures, Algorithms*, Springer (2006), to appear.
- [4] S. Elizalde and K. Woods, Bounds on the number of inference functions of a graphical model (2006), preprint.
- [5] M. Grötschel, L. Lovász, and A. Schrijver, *Geometric algorithms and combinatorial optimization*, Springer-Verlag, Berlin (1993).
- [6] C.G. Lekkerkerker, *Geometry of Numbers*, Wolters-Noordhoff, Groningen (1969).
- [7] J.E. Nymann, On the probability that k positive integers are relatively prime, *J. Number Theory* 4 (1972), 469–473.

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755
E-mail address: `sergi.elizalde@dartmouth.edu`

DEPARTMENT OF MATHEMATICS, OBERLIN COLLEGE, OBERLIN, OH 44074
E-mail address: `kevin.woods@oberlin.edu`